

AD\_\_\_\_\_

Award Number: DAMD17-98-1-8213

TITLE: Genetic Epidemiology of Mammographic Breast Density

PRINCIPAL INVESTIGATOR: Thomas A. Sellers, Ph.D.

CONTRACTING ORGANIZATION: Mayo Foundation  
Rochester, Minnesota 55905

REPORT DATE: October 2000

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;  
Distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20010521 071

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 074-0188	
<small>* Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 2000	3. REPORT TYPE AND DATES COVERED Annual (1 Oct 99 - 30 Sep 00)		
4. TITLE AND SUBTITLE Genetic Epidemiology of Mammographic Breast Density		5. FUNDING NUMBERS DAMD17-98-1-8213		
6. AUTHOR(S) Thomas A. Sellers, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Mayo Foundation Rochester, Minnesota 55905  E-MAIL: sellers.thomas@mayo.edu		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES This report contains colored photos				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; Distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words)  Mammographic breast density, has been shown to be a major risk factor for breast. We hypothesize that there is at least one major gene involved in the genetic variation of breast density. We propose to identify the chromosomal location for this gene(s) by linkage analysis. To do so, we have selected 23 large families providing the greatest information for linkage from a total of 426 breast cancer families in the Minnesota Breast Cancer Family Study. Mammograms and risk factor data were previously collected on these women through interview and mailed questionnaires. As of October 30, 2000, 384 samples from these families have been collected for analysis. We have also concentrated efforts on improving the breast density estimate. We have updated the subjective estimate of breast density to a computer-assisted estimate. Additionally, in the past year, we have performed several small studies to evaluate the reliability of this computer-estimate of density. The final year of the study will involve a genome screen for loci linked to the breast density trait, using a screening set of microsatellite markers that span the genome. The discovery of breast density genes could help in the identification of susceptible individuals to target prevention strategies.				
14. SUBJECT TERMS Breast Density, Breast Cancer			15. NUMBER OF PAGES 11	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)  
Prescribed by ANSI Std. Z39-18  
298-102

## Table of Contents

Cover.....	1
SF 298.....	2
Introduction.....	4
Body.....	5
Key Research Accomplishments.....	8
Reportable Outcomes.....	9
Conclusions.....	10
References.....	11
Appendices.....	n/a

## Introduction

The interindividual variability in breast tissue on mammographic images, as defined by several measures of mammographic breast density, has been shown to be a major risk factor for breast cancer with three to five-fold increases in risk associated with densities greater than 50% (Boyd, 1998). We have previously demonstrated a genetic component to the mammographic breast density trait (Pankow, 1997). Our hypothesis is that there is at least one major gene involved in the genetic variation of breast density, but there are no obvious candidate genes or chromosomal regions known with certainty. We now propose to identify the chromosomal location for this gene(s) by linkage analysis, as the first step in identifying the gene(s) that is responsible for the differences in breast density. In the body of this report, we will describe the progress to date on the search for the breast density gene(s). The discovery of breast density genes could help in the identification of susceptible individuals to target prevention strategies as well as provide insight into the development of breast cancer.

## Body

The goal of our study is to locate a chromosomal region for a gene for breast density. The major emphasis of this first phase of this linkage study has been to select informative families for linkage and collect DNA and trait information on these family members. We selected 23 families for linkage, based on simulation analyses that incorporated the trait information. We collected 384 blood samples on eligible women in these families who had mammogram and risk factor information. Our participation rate for the blood component of the study was 79%. The distribution of bloods per family is described in the table below.

Table 1. Distribution of bloods for 23 families in linkage analysis


Family No.	No. Bloods
018	11
042	9
055	15
086	16
115	29
165	39
198	8
263	10
285	22
316	14
400	14
406	16
418	15
433	8
438	31
464	12
498	17
504	10
506	19
540	16
549	27
588	18
589	8

The blood collection phase is now complete. As a result, the focus of the study will now shift to genotyping a panel of microsatellite markers for these 384 individuals. The panel consists of a screening set of microsatellite markers that are distributed across all autosomes and separated at an average distance of 10 centimorgans (ABI marker set). We approximate 6-8 months to complete the genotyping of these 23 families.


Also integral to any linkage analysis is a clearly defined trait. In our case, the trait of interest is mammographic breast density, which is defined as the percent of the breast that is composed of nonfatty tissue. However, there are several ways in which to measure percent density. Previously, in our segregation analysis, we used a subjective estimate by an experienced radiologist, which was shown to have high reliability and validity (compared to an MRI measure of the breast). However, we desired a more precise measure and have acquired software for a computer-assisted estimate of breast density developed by Martin Yaffe and colleagues at the University of Toronto (Yaffe, 1994). The procedure for setting thresholds on the mammographic image to arrive at a density estimate is pictorially described in Figure 1.

In the past year, we have performed studies to evaluate the computer-assisted estimate. We have trained a programmer to apply the algorithm and continually compute her intra and inter reliability estimates (comparing her estimates to those of Dr. Norman Boyd at the University of Toronto, an expert in breast density estimation). Her intra reader reliability is consistently above 90%, and her inter reader reliability with Dr. Norman Boyd on a set of 100 digitized images was 95%. Once the programmer has completed reading the mammograms for this study, we will compute the correlation between the subjective estimate of density and the computer-assisted estimate. If the correlation exceeds 90% then, we will perform our linkage analyses using the computer-assisted estimate of breast density rather than the subjective estimate.


Figure 1: Example of Percent Density Estimation



Digitized mammogram



Setting thresholds



Dense area highlighted

## Key Research Accomplishments

- Performed simulation analyses under nonparametric and parametric linkage to determine the most informative families for linkage analysis.
- Collected blood from 384 family members in 23 families with trait and risk factor information. Participation rate was 79%.
- Implemented a computer-assisted algorithm for estimation of percent mammographic breast density.
- Performing a comparison of the estimates from computer-assisted method with subjective estimates of density.
- Performed intra and inter reliability studies to evaluate the performance of the computer-assisted method.
- Currently are preparing to begin genotyping for the genome screen. Estimated completion time will be 6-8 months.



## Reportable Outcomes

Work as of 10/2000 has involved blood collection and DNA extraction for 384 participants. The only reportable outcome was our poster presentation at the DOD meeting last June.

Vachon CM, Thibodeau SN, Kulby VJ, Sellers TA. Genetic linkage analysis of mammographic breast density. Department of Defense Breast Cancer Research Program Meeting, June 2000.

## Conclusions

The majority of the past two years has involved the collection of blood samples as sources of DNA for the genome screen. Our participation has been above average, with 79% of women participating in the study. We have also devoted a large effort to developing a more adequate estimate of breast density. Having performed several studies on the performance of the computer-assisted method, we are confident that this measure will be appropriate for our linkage analysis. The final year of the study will involve the genotyping for the 384 women in these families. We will perform both parametric and nonparametric linkage analyses to hopefully arrive at a chromosomal location for a breast density gene.

## References

Pankow, J.S., Vachon, C.M., Kuni, C.C., King, R.A., Arnett, D.K., Grabrick, D.M., Rich, S.S., Anderson, V.E. and Sellers, T.A., Genetic analysis of mammographic breast density in adult women: evidence of a gene effect. *J. Natl. Cancer Inst.*, 89, 549-556 (1997).

Boyd, N.F., Lockwood, G.A., Byng, J.W., Tritchler, D.L. and Yaffe, M.J., Mammographic densities and breast cancer risk. *Cancer Epidemiol. Biomarkers Prev.*, 7, 1133-1144 (1998).

Byng JW, Boyd NF, Fishell E, Jong RA, Yaffe MJ. The quantitative analysis of mammographic densities. *Phys. Med. Biol.* 39,1629-1638 (1994).